

Effect of transplantation of allogeneic multipotent mesenchymal bone marrow stromal cells on regeneration of liver after extended hepatectomy (experimental study)

Titova A., Voskanyan S.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

© 2018 Human Stem Cell Institute. All rights reserved. Multipotent mesenchymal stromal cells can be a drug for treatment of acute liver failure. The purpose of this study was to assess the effect of multipotent mesenchymal stromal cells on the proliferative, mitotic activity of hepatocytes and morphometric criteria for the regeneration of liver parenchyma after 68 % partial hepatectomy in rats. The number of mitosis, PCNA positive cells, diameter of nucleus of hepatocytes, size of hepatocytes, nuclear-cytoplasmic ratio (NCR), the area of hepatic lobules were evaluated on 1, 2, 4, 14 days after surgery. The results of the study showed that intravenous administration of allogeneic mesenchymal multipotent stromal cells from bone marrow after extended hepatectomy increase the number of proliferating hepatocytes (PCNA positive cells), but does not affect the number of mitoses, the diameter of the nucleus, the area of hepatocytes, NCR and hepatic lobules. Probably, at this time, regeneration proceeds primarily through an intracellular mechanism.

<http://dx.doi.org/10.23868/201808025>

Keywords

Extended hepatectomy, Inferior vena cava, Multipotent mesenchymal stromal cells, Transplantation

References

- [1] Putignano A., Figorilli F., Alabsawy E. et al. Long-term outcome in patients with acute liver failure. *Liver Int.* 2018; 10.1111/liv.13914.
- [2] O'Grady J. Timing and benefit of liver transplantation in acute liver failure. *J Hepatol.* 2014; 60(3): 663-70.
- [3] Watt K.D., Pedersen RA, Kremers W.K. et al. Evolution of Causes and Risk Factors for Mortality Post Liver Transplant: Results of the NIDDK Long Term Follow-up Study. *American Journal transplantation* 2010; 10(6): 1420-7.
- [4] Sun X.E., Zhang X.Q., Liu M.M. Effect of bone marrow mesenchymal stem cells on the TGF- β 1/Smad signaling pathway of hepatic stellate. *Genetic and Molecular Research* 2015; 14: 8744-54.
- [5] Michalopoulos G.K., Marie C. Liver Regeneration. *DeFrances Science* 1997; 276: 60-6.
- [6] Liu W., Song F., Ren L.N. et al. The multiple functional roles of mesenchymal stem cells in participating in treating liver diseases. *Cell. Mol. Med.* 2015; 3: 511-20.

- [7] Cen P., Chen J., Hu C. et al. Noninvasive in-vivo tracing and imaging of transplanted stem cells for liver regeneration. *Stem Cell Research & Therapy* 2016; 7: 1-11.
- [8] Akhan E., Tuncel D., Akcali K.C. Nanoparticle Labeling of Bone Marrow-Derived Rat Mesenchymal Stem Cells: Their Use in Differentiation and Tracking. *Biomed. Res. Int.* 2015; 19: 511-20.
- [9] Liu T., Mu H., Shen Z. et al. Autologous adipose tissue-derived mesenchymal stem cells are involved in rat liver regeneration following repeat partial hepatectomy. *Mol. Med. Rep.* 2016; 13(3): 2053-9.
- [10] Yu J., Yin S., Zhang W. et al. Hypoxia preconditioned bone marrow mesenchymal stem cells promote liver regeneration in a rat massive hepatectomy model. *Stem Cell Research & Therapy* 2013; 4: 1-9.
- [11] Baretta G.A., Gama Filho O., Toderke E.L. et al. Effect of cyclosporine on liver regeneration in partial hepatectomized rats. *Acta Cir. Bras.* 2015; 30(1): 54-9.
- [12] Biondo-Simões M.L.P., Matias J.E.F., Montibeller G.R. et al. Effect of aging on liver regeneration in rats. *Acta Cirúrgica Brasileir* 2006; 1(4): 197-202.
- [13] D'Espressailles A., Dossi C., Intriago G. et al. Hormonal pretreatment preserves liver regenerative capacity and minimizes inflammation after partial hepatectomy. *Annals of hepatology* 2013; 12(6): 981-91.